

# Systematic Review on Quality of Life of Patients with Chronic Kidney Disease

Arpita J. Parekh and Dr.HN. Ravindra

**Abstract--- Background:** *QOL is an overall assessment of a person's well-being, which may include physical, emotional, and social dimensions, as well as stress level, sexual function, and self-perceived health status. End-stage renal failure is a chronic disease that exerts a great negative impact on patients' health-related QOL mainly due to the accompanied impairment or to the imposed limitations in almost all domains of their daily lives. Despite remarkable advances in the treatment of HD, the patients encounter certain physical, psychological, economic, and social problems which affect their QOL.*

**Objectives:** *This review aims to examine systematically the impact of patient navigator training on Quality Of Life of Patients with Chronic Kidney Disease undergoing haemodialysis to identify: (i) QOL of patients with CKD undergoing Hemodialysis (ii) QOL of patients with CKD undergoing Hemodialysis in terms of Client centred health education, Meditation & progressive muscle relaxation technique.*

**Methodology:** *Searches were conducted across 10 electronics data bases consistent keywords. Papers were screened by the title and abstract (n=733) and judged against pre-defined eligibility (n= 26). Twenty six papers were included in the review.*

**Results:** *Different patient navigator training approaches were included such as client centred health education to improve knowledge related to disease condition; meditation program which includes pranayama, progressive muscle relaxation technique and mindful meditation which may indirectly impact on quality of life (Physical health, Psychological, Social relationships & Environment). Patient navigator training may improve Quality Of Life of Patients with Chronic Kidney Disease undergoing haemodialysis.*

**Conclusion:** *Few studies reported effect on QOL (n=18), the lack of a clear theoretical framework resulted in ambiguous predictions about training and its effect. Methodological issues also constrain the strength of the evidence, including; small samples sizes; an absence of blinding of participants, outcome assessors; and lack of active control groups.*

**Keywords---** *QOL, Client Centred Health Education, Meditation, Progressive Muscle Relaxation Technique, Chronic Kidney Disease and Haemodialysis.*

---

## I. INTRODUCTION

The term **chronic kidney disease** applies to the process of continuing significant irreversible reduction in nephron number. The dispiriting term end-stage renal disease represents a stage of CKD where the accumulation of toxins, fluid, and electrolytes normally excreted by the kidneys results in the uremic syndrome. This syndrome leads

---

Arpita J. Parekh, PhD Scholar, Sumandeep Vidyapeeth Deemed to be University, Vadodara, Gujarat, India.  
E-mail: arpita.parekh03@gmail.com

Dr.HN. Ravindra, Professor, Sumandeep Nursing College, Sumandeep Vidyapeeth Deemed to be University, Vadodara, Gujarat, India.

to death unless the toxins are removed by renal replacement therapy, using dialysis or kidney transplantation.<sup>1</sup> chronic kidney disease is an increasing public health issue. Prevalence is estimated to be 8–16% worldwide<sup>2</sup>

The evaluation of **quality of life (QoL)** in chronic diseases is becoming more and more important.<sup>3</sup> The assessment of QoL helps in evaluating the quality of care and efficacy of medical intervention, improving clinical decision-making, and estimating health care needs of the community.<sup>4</sup> Treatment modalities for end-stage renal disease affect quality of life (QOL) of the patients. Moreover, health care personnel can use QoL scores to evaluate the effects of a specific disease on patients and specific treatment besides the assessment of progress of patients. The different interventions could provide benefits by improving patient adherence to medications, which could help slow the progression of kidney disease through better BP control. Furthermore, through patient education, rates of follow-up with health care providers could be improved.<sup>5</sup>

**Quality of life** is an important criterion that illustrates the effectiveness of health care, health level, and well-being. It is a multidimensional concept that includes ability, function, health, well-being, and psychological state, which is defined by the World Health Organization as values, goals, standards, and individual interests.<sup>6-8</sup> There is a relationship between diseases and quality of life. Quality of life can have a direct impact on physical performance, emotional, and physical problems, fatigue, mental health, social performance, physical pain, and general health.<sup>7-8</sup> Therefore, knowledge about chronic diseases, especially chronic kidney diseases (CKD) is very important in the evolution of patients' health problems.

The role of nurses in imparting **education** about a proper diet to patients on **HD** is crucial. Patients on HD experience decreased quality of life (QoL) and significantly higher rates of malnutrition, inflammation, hospitalization and mortality compared with the normal population.<sup>9</sup> **Health educational** interventions which provide information about the nature of illness, exercise and relevant coping skills have also been used and found to enhance the physical and psychosocial well-being of patients with CKD<sup>10</sup> Therefore interventions which are able to enhance physical and mental functions can reduce hospitalization and mortality rates in CKD patients. The other interventions which have been used in this population have focused on education, social support, and self-care and these have been shown to raise coping skills among people on dialysis.<sup>11</sup>

**Relaxation therapy** techniques and mindfulness meditation are clinical intervention tools that have demonstrated benefits associated with the reduction of negative psychological states and the enhancement of positive states of mind critical to the alleviation of physical and emotional distress<sup>12</sup>.

**Meditation** has been viewed as a potential intervention to decrease stress and anxiety and improve quality of life has seen an increase in popularity over the past decade. Meditation is an example of a mind-body intervention as it helps to influence the mind to adapt to the body's physical symptoms through mechanisms of the parasympathetic nervous system and the decrease in stress hormone levels. Other mind-body interventions include relaxation techniques, hypnosis, yoga, psychotherapy, guided imagery, prayer, dance, music, and art therapy.<sup>13</sup> Pranayam-Yoga program is safe and effective measure as adjuvant therapy to conventional treatment modalities in reducing blood pressure, improving renal function, decreasing the need for dialysis, and improving QOL in patients with CKD. As

CKD had a chronic course whether the results of our short-term study can be extrapolated to long-term benefits is yet to be explored.<sup>14</sup>

### **Objectives**

This review aims to examine systematically Quality Of Life of Patients with Chronic Kidney Disease undergoing haemodialysis to identify:

- (i) QOL of patients with CKD undergoing Hemodialysis
- (ii) QOL of patients with CKD undergoing Hemodialysis in terms of Client centred health education, Meditation & progressive muscle relaxation technique.

### **Review Method**

Data searches: A literature search was performed in the following electronic database: PubMed, CINAHL plus, Google Scholar, Research gate, Science direct, EBSCOhost, PsycINFO, SCOPUS, Web of Science, Shodhganga were used to search the literature for all publications from 2009 to 2019 (last 10 years).

Keywords used - “quality of life” or “health related quality of life “ and chronic kidney disease\* AND (training or intervention \* or treatment or therapy or program\*)

## **II. SEARCH STRATEGY**

**Inclusion criteria: “PICO” framework**

**Inclusion and exclusion criteria for article selection**

**Inclusion criteria: “PICO” framework**

**Population**– Studies must have been conducted with children aged above 18 years

**Intervention** –Studies must have implemented any intervention that to improve quality of life with chronic kidney disease patients.

**Comparisons** –Studies must have a randomized controlled, quasi-experimental, or single case experimental design.

**Outcomes** –Studies must have at least one pre-and post-intervention measures of quality of life

Primary outcome- impact on quality of life

Secondary outcomes –improvement in QOL (physical health, psychological health, social and environmental domains)

The research paper only which directly belongs to quality of life with chronic kidney disease patient. The paper which is easily accessible online and full text available.

The studies which are carried out in English language.

### **Exclusion criteria**

The study concern to chronic kidney disease patient receiving haemodialysis

The studies which are duplicate and result are ambiguous.


The research study which is published in without ISSN No journals.

The research studies which are not available on journal database.

The research studies in which only abstract is available.

The studies which is published in local language.

### Study Selection



**Figure 1- Flow diagram of search strategy with reasons of exclusion.**

To start with from below listed search engines total 733 articles were received; from that 297 articles were excluded bases on exclusion criteria. So total retrieved articles were 436 among all 89 duplicate articles, 94 review articles and 253 abstracts were excluded. Final retrieved articles were 94; among them 68 full articles were excluded based on inclusion criteria. Finally, 26 were included in the review.

### III. RESULTS

All included studies have been classified into four intervention types –

1. **Quality of life among CKD patients undergoing haemodialysis (n=8):** Eight studies reviews here investigated the existing quality of life of patients undergoing haemodialysis with chronic kidney disease.
2. **QOL of patients undergoing haemodialysis in terms of Client centred health education, Meditation & progressive muscle relaxation technique (n=18):** Interventions divided into below 4 categories.
  - 1) **Client centred health education (n=4):** Four interventions related to health education on disease process, prognosis, nutrition and diet.
  - 2) **Meditation (n=6):** Wide range of meditation and yoga showed effectiveness on improvement of quality of life in all domains. It has been suggested systematic regular meditation may benefit for CKD patients to improve quality of life.

- 3) **Progressive Muscle relaxation Technique (n=5):** Five studies suggested progressive muscle relaxation technique in organized manner can create impact on physical health of an individual.
- 4) **Patient navigator training (n=3):** Three interventions involved repeated combine practice may produce progressive effect on quality of life. Multidimensional combination of different intervention like health education, regular meditation activities and relaxation techniques may create positive effect on quality of life of patients with chronic kidney disease.

Table 1: Summary of Characteristics of all Included Studies in Review (n=26)

| Article/<br>Ref No                                      | Study Title                                                                              | Country              | Year of<br>study             | Sample                                                     | Study Type                                                                     | Methodological aspects                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | Findings                                                                                                                                                                                                                          |
|---------------------------------------------------------|------------------------------------------------------------------------------------------|----------------------|------------------------------|------------------------------------------------------------|--------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>QOL of patients with CKD undergoing Hemodialysis</b> |                                                                                          |                      |                              |                                                            |                                                                                |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |                                                                                                                                                                                                                                   |
| 1/15                                                    | Predictors of Quality of Life in Haemodialysis Patients                                  | Ismailia, Egypt      | January 2007 to January 2008 | 100 HD patients<br>53: Males<br>47: Females                | A cross-sectional descriptive design                                           | SF-36 and KDQoL-SF forms covering six domains of QoL, namely physical, emotional, social, illness impact, medical and financial satisfaction, and overall general health                                                                                                                                                                                                                                                                                                                                                         | Quality of life is reduces in all the health domain of haemodialysis patient. Aged male, unemployment and duration of dialysis adversely affect the quality of life score. A novel approach needed to influence patient outcomes. |
| 2/16                                                    | An assessment of the quality of life in hemodialysis patients                            | Mysore , India       | October 2008                 | 75 : HD patients                                           | A cross-sectional study                                                        | The WHOQOL-BREF questionnaire was used to assess the quality of life. The quality of life of hemodialysis patients was found to be significantly impaired ( $P < 0.05$ ) in comparison to healthy individuals of the general population, particularly with respect to the physical, psychological, and social relationship domains. In comparison to the quality of life of renal transplant patients, the quality of life of haemodialysis patients was significantly ( $P < 0.05$ ) lower in all the four WHOQOL-BREF domains. | The quality of life of hemodialysis patients was found to be considerably impaired when compared to that of healthy individuals of the general population as well as of renal transplant patients.                                |
| 3/17                                                    | Dialysis-Related Factors Affecting Quality of Life in Patients on haemodialysis          | Lahore, Pakistan     | 2011                         | 175: HD patients                                           | Cross-sectional study                                                          | Data collected on patient on maintenance haemodialysis for more than 3 months at 3 dialysis centres of Lahore. The QOL index measured using the World Health Organization QOL questionnaire, with higher scores corresponding to better QOL of patients.                                                                                                                                                                                                                                                                         | QOL of haemodialysis patients is poor as compared to caregivers of the patients, especially that of diabetics. Also, duration of dialysis had a reverse correlation with QOL.                                                     |
| 4/18                                                    | Assessment of Quality of Life in Patients on Haemodialysis and the Impact of Counselling | Kochi, Kerala, India | 2012                         | 50 patients<br>25: Experimental Group<br>25: Control Group | A hospital-oriented prospective, longitudinal, observational comparative study | QOL of patients on haemodialysis by using the World Health Organization Quality of Life assessment scale for six months in the nephrology department of a tertiary care hospital. Patients who were                                                                                                                                                                                                                                                                                                                              | There is an increase in score in all the four domains (physical, psychological, environmental and Social) among the test group when compared with the                                                                             |

|                                                                                                     |                                                                                                        |                             |               |                                                                 |                                 |                                                                                                                                                   |                                                                                                                                                                                                                                                                                                        |
|-----------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------|-----------------------------|---------------|-----------------------------------------------------------------|---------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                                                     |                                                                                                        |                             |               |                                                                 |                                 | receiving HD regularly and aged between 20 and 80 years were included in the study.                                                               | control group. Also, found that the Psychological domain shows significant increase in score compared with others. Findings demonstrate that patient counselling plays an important role in improving the QOL.                                                                                         |
| 5/19                                                                                                | Assessment of Quality of Life In Chronic Kidney Disease Patients                                       | Chennai, Tamil Nadu, India. | July 2014     | 50 patients                                                     | Observational and prospective   | Observed by using kidney disease QOL short form TM (KDQOL-SFTM) questionnaire.                                                                    | Findings revealed that ESRD patients have a poor QOL and most the affected domain is PH, hence measuring and monitoring These aspects of QOL could lead to a more patient-centered care and improve the health and well-being among patients with chronic renal failure.                               |
| 6/20                                                                                                | Assessment of health-related quality of life of hemodialysis patients                                  | Benha, Egypt                | December 2015 | 228:HD patients                                                 | A cross-sectional study         | Data collected by questionnaire that included demographic, social, and medical questions. KDQOL SF (36) as a tool for assessment of HRQOL.        | HRQOL in the hemodialysis patients relatively low. The most important comorbidities affecting quality of life were anemia, HCV infection, and diabetes.                                                                                                                                                |
| 7/21                                                                                                | Quality of Life of Patients Undergoing Hemodialysis                                                    | Tamil Nadu, India           | January 2017  | 130 : HD patients                                               | Cross-sectional research design | KDQOL - short form version 1.3 scale used for the study to collect the demographic variable and data.                                             | Concluded that patients on HD not having adequate QOL in all domains except patient satisfaction due to changes in the physiological, chemical changes occur in the kidney.                                                                                                                            |
| 8/22                                                                                                | The impact on quality of life of dialysis patients with renal insufficiency                            | Poland                      | 2018          | 140 dialysis patients<br>100 : HD<br>40 : PD                    | Observational and prospective   | The questionnaire used for data collection by Kidney Disease and Quality of Life Short Form version 1.2 (KDQOL – SF 1.2)                          | Patients receiving peritoneal dialysis assessed their QoL in its different dimensions as much higher than patients receiving hemodialysis. The data also conclude that will to live was more highly assessed by patients receiving peritoneal dialysis as compared to patients receiving hemodialysis. |
| <b>QOL of patients with CKD undergoing Hemodialysis in terms of Client centred health education</b> |                                                                                                        |                             |               |                                                                 |                                 |                                                                                                                                                   |                                                                                                                                                                                                                                                                                                        |
| 9/23                                                                                                | The Impact of Education on Nutrition on the Quality of Life in Patients on Hemodialysis: A Comparative | Urmia, Iran                 | 2012          | 70 : HD patients<br>35: Experimental Group<br>35: Control Group | A quasi experimental study      | Seventy patients on maintenance haemodialysis divided into two groups; both groups requested to fill in the validated SF-36 questionnaire on QoL. | The QoL considerably diminished in HD patients. One of the methods for this is education about their nutritional program, which can be used for other                                                                                                                                                  |

|                                                                                |                                                                                                                           |                      |              |                                                             |                                          |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |                                                                                                                                                                                                                                                    |
|--------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------|----------------------|--------------|-------------------------------------------------------------|------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                                | Study from Teaching Hospitals                                                                                             |                      |              |                                                             |                                          |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             | chronic diseases too.                                                                                                                                                                                                                              |
| 10/24                                                                          | The effect of education of health-promoting behaviours on lifestyle in haemodialysis patients                             | Iran                 | 2014-15      | 70: HD patients                                             | Quasi-experimental study                 | Haemodialysis patients in two groups: control [35] and experimental [35]. For groups of six 30-minute sessions on an individual basis during haemodialysis, held over three weeks in a row.                                                                                                                                                                                                                                                                                                                 | Teaching with an emphasis on health-promoting behaviours, haemodialysis patients was improved lifestyle.                                                                                                                                           |
| 11/25                                                                          | Effectiveness of education and exercise on quality of life among patients undergoing hemodialysis                         | Muscat, Oman         | October 2017 | 150 HD patients<br>75: in intervention<br>75: control group | A randomized controlled trial            | Laptop assisted interactive health education and supervised the exercise during dialysis given in vernacular language for 25 mins to the experimental group. Data collected through Research and Development health-related kidney disease QOL (KDQOL) questionnaire                                                                                                                                                                                                                                        | The education and exercise intervention have a positive effect on the physical and mental health well being of patients with chronic kidney disease.                                                                                               |
| 12/26                                                                          | The effect of an educational program on quality of life in patients undergoing hemodialysis                               | Western Saudi Arabia | 2019         | 100 : HD patients                                           | A randomized interventional study design | 100 HD patients recruited and randomly assigned to control group(50) and interventional group (50) from four different hospital data collected by HRQOL SF-36. Health education (twice a week, total duration of 8 weeks, 30 minutes each session) sessions which focus on an introduction to chronic renal disease, explaining HD, how it works, and why it is important to receive and to comply to HD conducted to the participants.                                                                     | The scores of HRQOL for HD patients were low. The educational program had significant positive impact on all health-related quality of life parameters.                                                                                            |
| <b>QOL of patients with CKD undergoing Hemodialysis in terms of Meditation</b> |                                                                                                                           |                      |              |                                                             |                                          |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |                                                                                                                                                                                                                                                    |
| 13/27                                                                          | Effects of 6 months yoga program on renal functions and quality of life in patients suffering from chronic kidney disease | Meerut, India        | 2009-2010    | 54 patients                                                 | Prospective study                        | In experimental group (25) trained to perform yogic asanas for at least 5 days/week for 40-60 mins/day. Control group (25) on conventional therapy. Both group assess by quality of life (QOL) indicators with 6 months follow up.                                                                                                                                                                                                                                                                          | Yoga program is safe and effective as an adjuvant therapy in improving renal functions and QOL of CKD patients.                                                                                                                                    |
| 14/28                                                                          | Role of Yoga in Chronic Kidney Disease:                                                                                   | Bangalore, India     | 2014         | A Hypothetical Review                                       | A Hypothetical Review                    | Different studies suggested that there is a strong association of oxidative stress, chronic inflammation and psychological stress with CKD. In several studies, it has been reported that yoga has significant role in the management of non-communicable diseases like diabetes, hypertension, coronary heart diseases etc. Regular yoga practice can help control sugar levels in diabetics, blood pressure in hypertensives and reduce the risk of cardiac complications in patients with heart diseases | Review suggested that yoga has promising role in the primary and secondary management of CKD as an adjuvant. These researches and based on this present a yoga module useful in CKD along with necessary precautions to be taken while doing yoga. |
| 15/29                                                                          | Improving                                                                                                                 | CA,                  | 2017         | A                                                           | A systematic                             | Meditation, as part of more                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 | The studies have                                                                                                                                                                                                                                   |

|                                                                                                   |                                                                                                                  |        |      |                         |                                      |                                                                                                                                                                                                                                                                                              |                                                                                                                                                                                                                                                                                                                                  |
|---------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------|--------|------|-------------------------|--------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                                                   | wellbeing in patients undergoing dialysis: Can Meditation Help?                                                  | USA.   |      | systematic review Study | review Study                         | rigorous and physically active programs, has been shown to be useful in dialysis contexts. Likewise, meditation alone may even benefit cognitive impairment in end-stage kidney disease patients.                                                                                            | shown varied results, though most have reported positive effects of meditation on measures such as anxiety, stress, depression, sleep disorders, quality of life and improve experience of undergoing dialysis. "Can Meditation Help the Wellbeing of Dialysis Patients?" the systematic review response is an optimistic "yes." |
| 16/30                                                                                             | Mindfulness Meditation Practice During Hemodialysis                                                              | U.S.A  | 2017 | 50: HD patients         | Randomized control trial             | In experimental group Mindfulness meditation practice based on relaxation, well-being promotion, meditation practices and positive psychology principles. The program is being evaluated in haemodialysis session for 12 weeks, for 15 -25 minutes, 3times/week.                             | Improvement of symptoms of depression, stress, quality of life, sleep disorders, biochemical parameters resulting from better adherence to treatment and Improvement of symptoms of quality of life                                                                                                                              |
| 17/31                                                                                             | Brief Mindfulness Meditation for Depression and Anxiety Symptoms in Patients Undergoing Hemodialysis             | Canada | 2017 | 41:HD patients          | Randomized control trial             | 41 patients; 21 experimental and 20 controls. Experimental group received 8 week, individual chair side meditation for 10-15 minutes 3times/week during haemodialysis.                                                                                                                       | Mindfulness meditation appears to be feasible and well tolerated in patients on haemodialysis with anxiety and depression symptoms                                                                                                                                                                                               |
| 18/32                                                                                             | The effect of mindfulness program on general health of patients undergoing hemodialysis                          | Iran   | 2018 | 60:HD patients          | Clinical trial (IRCT2017053134263N1) | In experimental group received 8 sessions of mindfulness training and education session. The control group received only 8 sessions of education in relation to ESRD and HD. Immediately after and 1 Month after the intervention, the General Health Questionnaire assessed by both groups. | Mindfulness is effective in reducing physical and anxiety symptoms, sleep disorder, social dysfunction, and depression symptoms. Hence, the use of mindfulness as a complementary treatment can improve the general health level in these patients.                                                                              |
| <b>QOL of patients with CKD undergoing Hemodialysis in terms of progressive muscle relaxation</b> |                                                                                                                  |        |      |                         |                                      |                                                                                                                                                                                                                                                                                              |                                                                                                                                                                                                                                                                                                                                  |
| 19/33                                                                                             | The effect of progressive muscle relaxation training on anxiety levels and quality of life in dialysis patients. | Turkey | 2006 | 46: Dialysis patients   | Quasi-experimental study             | Sample collected by patient recognition form, state and trait anxiety inventory and QoL – index for dialysis patients. Progressive muscle relaxation technique administered for 6 weeks.                                                                                                     | PMRT for dialysis patients helps decrease state- and trait-anxiety levels and has a positive impact on QoL.                                                                                                                                                                                                                      |
| 20/34                                                                                             | The effects of an interventional program based on self-care model on                                             | Iran   | 2012 | 32 : HD patients        | A quasi-experimental study           | The self-care: partnership care model used for intervention. This model consists of four stages:<br>(1) Motivating,<br>(2) Preparing,                                                                                                                                                        | Implementing the self-care model increased the quality of life of haemodialysis patients. It is recommended for the                                                                                                                                                                                                              |

|                                                                                                |                                                                                                                                                                    |                   |                   |                           |                              |                                                                                                                                                                                                                                                                    |                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------|-------------------|---------------------------|------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                                                | health- related quality of life outcomes in hemodialysis patients                                                                                                  |                   |                   |                           |                              | (3) Involving, and<br>(4) Evaluating.<br>Required data collected by the Short Form- 36 (SF- 36) standard questionnaire and a researcher- made questionnaire.                                                                                                       | use of this model in haemodialysis patients.                                                                                                                                                                                                                                                                                                                                                                                                  |
| 21/35                                                                                          | Effect of Progressive Muscle Relaxation and Aerobic Exercise on Anxiety, Sleep Quality, and Fatigue in Patients with Chronic Renal Failure Undergoing Hemodialysis | Iran              | 2016              | 100:HD patients           | Double-blind clinical trial  | Progressive muscle relaxation technique, aerobic exercise and control given for 60 days. Questionnaires of anxiety, sleep quality and fatigue completed by participants before and after interventions.                                                            | PMR compared to aerobic exercise in improving the symptoms of anxiety, sleep disorders, quality of life and fatigue in haemodialysis patients                                                                                                                                                                                                                                                                                                 |
| 22/36                                                                                          | Effectiveness of Intra-dialytic Exercise on Dialysis Adequacy, Physiological Parameters, Biochemical Markers and Quality of Life – A Pilot Study                   | Tamil Nadu, India | March to May 2017 | 20 : HD Patients          | Experimental research design | Experimental group received intra-dialytic exercise session for 3 times/week by using bicycle ergometer for 12 weeks during the first 2 h of HD besides receiving routine care compared to the control group                                                       | The prescribed intra-dialytic exercise intervention resulted in significant improvement in Kt/V, serum creatinine, blood urea, serum potassium, phosphorous, and quality of life with no adverse effects. This exercise program is safe complementary intervention and do not cost the patient extra time.                                                                                                                                    |
| 23/37                                                                                          | Effect of Multidimensional Educational Interventions Among Dialysis Patients                                                                                       | Karnataka India   | 2018              | A systematic review Study | A systematic review Study    | Multidimensional Educational Interventions : Effectiveness of Nutritional Intervention and Fluid Restriction on Patient Outcomes, Exercise Counselling and Patient Outcomes, Medication Counselling and Patient Outcomes, Quality of Life and Self-Care Management | Individual discipline proven educational interventions effective among dialysis patients to improve quality of life, self-care management, knowledge, biochemical parameters and therapeutic adherence. It is recommended that multidisciplinary approach covering dietary, nursing, primary physician, physical therapist and counsellor using coordinated strategies can aim and prove to improve the patient's outcomes to larger extents. |
| <b>QOL of patients with CKD undergoing Hemodialysis in terms of Patient Navigator Training</b> |                                                                                                                                                                    |                   |                   |                           |                              |                                                                                                                                                                                                                                                                    |                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| 24/38                                                                                          | Development of a chronic kidney disease patient navigator                                                                                                          | U.S.A             | 2012-13           | 2 : CKD Patients          | Randomized control trial     | CKD Patient navigator program included General patient navigator training, CKD education and EHR training in 5 institutions awarded a                                                                                                                              | <b>CKD patient navigator program CKD navigator hiring:</b> The appropriate personal and computer skills                                                                                                                                                                                                                                                                                                                                       |

|       |                                                                                                          |        |      |                           |                                             |                                                                                                                                                                                                                                                                                                                                                                                                                                  |                                                                                                                                                                                                                                                                                                                                                       |
|-------|----------------------------------------------------------------------------------------------------------|--------|------|---------------------------|---------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|       | program                                                                                                  |        |      |                           |                                             | grant to pilottranslational CKD interventions. CKD education provided by nurse practitioner during dialysis in 4 sessions 1 hour in each session about basic care of fistula and disease process.                                                                                                                                                                                                                                | outlined to meet the needs of the program<br><b>CKD patient navigator training:</b><br>Aim of the observation to make the patient navigator aware of the complexity of CKD, chronic disease management, and patients themselves so they may provide guidance and work with the patient and team to achieve the best possible outcome for the patient. |
| 25/39 | Pragmatic Randomized, Controlled Trial of Patient Navigators and Enhanced Personal Health Records in CKD | U.S.A. | 2017 | 209 patients              | 2*2 factorial design into four-study groups | Randomly in a 2 * 2 factorial design into four-study groups:<br>(1) Enhanced personal health record only,<br>(2) Patient navigator only,<br>(3) Both, and<br>(4) Usual care (control) group.<br>Primaryoutcome measure was the change in eGFR over a 2-year follow-up period.<br>Secondary outcome measures included acquisition of appropriate CKD-related laboratory measures, specialty referrals, and hospitalization rates. | Patient navigator program enhanced personal health record for the CKD population. Larger and long-term studies along with cost-effectiveness analyses are needed to evaluate the role of patient navigators and patient education through an enhanced personal health record in those with CKD.                                                       |
| 26/40 | Patient navigators for people with chronic disease: A systematic review                                  | Canada | 2017 | A systematic review Study | A systematic review Study                   | Searched MEDLINE, EMBASE, CENTRAL, CINAHL, PsycINFO, and Social Work Abstracts from inception to August 23, 2017. Also searched the reference lists of included articles.                                                                                                                                                                                                                                                        | Findings indicate that patient navigator programs improve processes of care, although few studies assessed patient experience, clinical outcomes or costs.                                                                                                                                                                                            |

#### IV. DISCUSSION

The aim of this review was to examine systematically the impact of Patient navigator training on Quality of Life of CKD patients undergoing hemodialysis

The first objective was to assess the Quality of Life of CKD patients undergoing hemodialysis. A systematic search of 8 studies encompassing a range of quality of life

Many of the studies clearly suggested that quality of life is compromised among patients of chronic kidney disease undergoing hemodialysis. In particular, quality of life of patients undergoing hemodialysis is more affected than undergoing peritoneal dialysis.

The second objective was to identify Impact of patient navigator training in terms of Client centred health education & Meditation on QOL. A systematic search of 18 studies in regards to health education 4 studies,

meditation 6 studies, progressive muscle relaxation technique 5 studies and patient navigator training 3 studies. Preliminary evidence suggesting that health education specific to the disease process improves their understanding and their vision towards the disease. A number of interventions reviewed those significant benefits by systematic meditation practice in improvement in all domains of quality of life. The effective progressive muscle exercise improves muscle cramp during dialysis procedure which indirectly improves physical health- domain of quality of life. Collectively all intervention together make one program like patient navigator training to improve quality of life of patients with CKD undergoing hemodialysis

## V. CONCLUSION

Diverse intervention applied within CKD patients undergoing hemodialysis in everyday context have produces improvements in their quality of life and have potential to produce near far effects. It's a challenging for patients with CKD undergoing hemodialysis. There are numbers of vital interventions combination can be highly effective to improve QOL of CKD patients. Client centered health education; meditation and progressive muscle relaxation technique combining as a one interventional program which may directly and indirectly create an impact on quality of life beneficial.

## VI. LIMITATIONS

Several methodological factors may limit confidence in the findings of this review including;

The risk of bias in individual studies due to the frequent lack of control of confounding variables,

Small sample sizes and Absence of blinding of participants and outcome assessors.

Lack of active control group

Absence of a clear theoretical framework in many included studies

## VII. RECOMMENDATION

- Future studies need to be more rigorous in study methodology including active control group and randomized controlled designs.

## REFERENCES

- [1] A. S. Fauci, D. L. Kasper, D. L. Longo. Harrison's Internal Medicine. 17th edition. First published: 29 December 2008
- [2] Prof Vivekanand Jha DM, Chronic kidney disease: global dimension and perspectives. *The Lancet Volume* 382, Issue 9888, 20–26 July 2013, Pages 260-272.
- [3] Rebollo P, Ortega F, Ortega T, Valdés C, García-Mendoza M, Gómez E. Spanish validation of the "Kidney Transplant Questionnaire": A useful instrument for assessing health related quality of life in kidney transplant patients. *Health Qual Life Outcomes* 2003; 1: 56.
- [4] Parekh, S. G., D. K. Ghosh, and S. R. Patel. "On frailty models for kidney infection data with exponential baseline distribution." *International Journal of Applied Mathematics & Statistical Sciences (IJAMSS)* 4.5 (2015): 31-40.
- [5] Ware JE, Brook RH, Davies-Avery. A: choosing measures of health status for individuals in general populations. *Am Public Health* 1981; 71:620-5.

- [6] BaharehGhiasi, Dian Sarokhani, Ali Hasanpour Dehkordi, Kourosh Sayehmiri , Mohammad Hossein Heidari et al. Quality of Life of patients with chronic kidney disease in Iran: Systematic Review and Meta-analysis. *Indian Journal of Palliative care*. Volume 24. Issue 1; 104-111.
- [7] Alam, M. SHABBIR, et al. "Effect of khat (*Catha edulis*) consumption on the functions of liver, kidney and lipid profile in male population of Jazan Region of Kingdom of Saudi Arabia." *Inter J Applied Natural Sci* 3.2 (2014): 9-14.
- [8] Unruh ML, Weisbord SD, Kimmel PL. Health-related quality of life in nephrology research and clinical practice. *Semin Dial* 2005; 18: 82-90.
- [9] BaharehGhiasi, Dian Sarokhani, Ali Hasanpour Dehkordi, Kourosh Sayehmiri, Mohammad Hossein Heidari et al. Quality of Life of patients with chronic kidney disease in Iran: Systematic Review and Meta-analysis. *Indian Journal of Palliative care*. Volume 24. Issue 1; 104-111.
- [10] Sharma, Surinder K., Indira R. Samal, and SOUMYA NP. "Urinary Stones in Southern India: Biochemical Analysis and its Clinical Implications." *International Journal of General Medicine and Pharmacy (IJGMP)* 4.1 (2015): 93-100.
- [11] Hasanpour-Dehkordi A, Khaledi-Far A, Khaledi-Far B, Salehi-Tali S. The effect of family training and support on the quality of life and cost of hospital readmissions in congestive heart failure patients in Iran. *Appl Nurs Res* 2016;31:165-9.
- [12] Nader Aghakhani. The impact of education on nutrition on the quality of life in patients on hemodialysis: a comparative study from teaching hospitals. *Saudi J Kidney Dis Transpl*. 2012;23(1):26-30.
- [13] Kimmel PL, Patel SS. Quality of life in patients with chronic kidney disease: focus on end-stage renal disease treated with hemodialysis. *Semin Nephrol*. Elsevier; 2006.
- [14] Karolich RL, Ford JP. Meaning, comprehension, and manageability of end-stage renal disease in older adults living with long-term hemodialysis. *Soc Work Health Care*. 2010;49(1):19-37
- [15] Gary Petingola. Relaxation Therapy and Mindfulness Meditation One-Day Workshop for Nephrology Social Workers: Exploring the Impact of This Training on Professional Practice. *National Kidney Foundation Journal of Nephrology Social Work*.
- [16] Paul N Bennett, Improving wellbeing in patients undergoing dialysis: Can meditation help Seminars in Dialysis. 2017;1-6. [wileyonlinelibrary.com/journal/sdi](http://wileyonlinelibrary.com/journal/sdi) ©2017 Wiley Periodicals, Inc.
- [17] Athab, Ahmed, Nabeel KM Ali, and Luma T Ahmed. "C-Reactive Protein and Renal Function Tests in Chronic Renal Failure Patients on Hemodialysis and Kidney Transplantation." *International Journal of Medicine and Pharmaceutical Science (IJMPS) ISSN (P)* (2016): 2250-0049.
- [18] Rajendra Kumar Pandey. Effects of 6 months yoga program on renal functions and quality of life in patients suffering from chronic kidney disease. *International journal of Yoga*. Year: 2017. Volume: 10. Issu : 1. Page : 3-8.
- [19] McBrien KA, Ivers N, Barnieh L, Bailey JJ, Lorenzetti DL, Nicholas D, et al. Patient navigators for people with chronic disease: A systematic review. *PLoS One*. 2018; 13(2): 1-23.
- [20] Obunwo, C., and J. Konne. "Measurement of levels of potassium bromate and some heavy metals in bread samples produced in Port Harcourt metropolis, Nigeria." *International Journal of Applied and Natural Sciences* 3.3 (2014): 1-4.
- [21] Dzikova S, Selim G, Gelev S, Trajceska L, Pushevski V, Sikole A. of Kidney Diseases and Transplantation Original Article Ultrasound Predictors of Compensated Liver Cirrhosis in Hemodialysis Patients with Hepatitis C. *Saudi J Kidney Dis Transplant [Internet]*. 2013; 24(1):30-5.
- [22] Parthasarathi G, Narahari M, Gurudev K, Sathvik B. An assessment of the quality of life in hemodialysis patients using the WHOQOL-BREF questionnaire. *Indian J Nephrol*. 2008; 18(4): 141.
- [23] Anees M, Hameed F, Mumtaz A, Ibrahim M, Khan MNS. Dialysis-related factors affecting quality of life in patients on hemodialysis. *Iran J Kidney Dis*. 2011; 5(1): 9-14.
- [24] Abraham S, Venu A, Ramachandran A, Chandran PM, Raman S. Assessment of quality of life in patients on hemodialysis and the impact of counseling. *Saudi J Kidney Dis Transpl*. 2012;23(5):953-7.
- [25] Murali R, Sathyanarayana D, Muthusethupathy M. Assessment of quality of life in chronic kidney disease patients using the kidney disease quality of life-short formtm questionnaire in indian population: A community based study. *Asian J Pharm Clin Res*. 2015; 8(1): 271-4.
- [26] Farahat HG, Marey HMG, Badawi NM, Allam HK, Issa MMM. Prevalence of occult refractive errors in primary school students. *Menoufia Med J*. 2017; 31(1): 267-72.
- [27] Thenmozhi P. Quality of life of patients undergoing hemodialysis. *Asian J Pharm Clin Res*. 2018; 11(4): 219-23.
- [28] Dąbrowska-Bender M, Dykowska G, Żuk W, Milewska M, Staniszevska A. The impact on quality of life

- of dialysis patients with renal insufficiency. *Patient Prefer Adherence*. 2018;12:577–83.
- [29] Esfehiani AJ, Dashti S. The impact of education on nutrition on the quality of life in patients on hemodialysis: a comparative study from teaching hospitals. *Saudi J Kidney Dis Transpl*. 2013; 24(1): 130.
- [30] Arani MD, Taghadosi M, Gilasi HR. The effect of education based on BASNEF model on lifestyle in patients with hypertension. *Iran Red Crescent Med J*. 2017;19(11):214–20.
- [31] Lazarus ER. Effectiveness of education and exercise on quality of life among patients undergoing hemodialysis. *Clin Epidemiol Glob Heal [Internet]*. 2019;7(3):402–8.
- [32] Bakarman MA, Felimban MK, Atta MM, Butt NS. The effect of an educational program on quality of life in patients undergoing hemodialysis in western Saudi Arabia. *Saudi Med J*. 2019; 40(1): 66–71.
- [33] Rk P. Users online: 87 Ahead of print | Login. 2020;1–7.
- [34] GM K. Role of Yoga in Chronic Kidney Disease: A Hypothetical Review. *J Nephrol Ther*. 2014;04(03).
- [35] Bennett PN, Ngo T, Kalife C, Schiller B. Improving wellbeing in patients undergoing dialysis: *Can meditation help? Semin Dial*. 2018;31(1):59–64.
- [36] Thomas Z, Novak M, Platas SGT, Gautier M, Holgin AP, Fox R, et al. Brief mindfulness meditation for depression and anxiety symptoms in patients undergoing hemodialysis a pilot feasibility study. *Clin J Am Soc Nephrol*. 2017; 12(12): 2008–15.
- [37] NCT03162770. Mindfulness Meditation Practice During Hemodialysis. <https://clinicaltrials.gov/show/NCT03162770> [Internet]. 2017;1–8.
- [38] Mohebi S, Parham M, Sharifirad G, Gharlipour Z. Social Support and Self- Care Behavior Study. 2018;1–6.
- [39] Momo Lokko, Christiana Naa, and Francis Lokko. "Clinical Presentation and Management of Human Ebola Virus Disease in Sub-Saharan Africa." *International Journal of Medicine and Pharmaceutical Science (IJMPS) ISSN (P)* (2016): 2250-0049.
- [40] Yildirim YK, Fadiloglu C. The effect of progressive muscle relaxation training on anxiety levels and quality of life in dialysis patients. *EDTNA-ERCA J*. 2006;32(2):86–8.
- [41] Ghavidel F, Mohammadzadeh S, Ravangard R, Bahadori M. The effects of an interventional program based on self-care model on health-related quality of life outcomes in hemodialysis patients. *J Educ Health Promot*. 2014; 3(1):110.
- [42] Amini E, Goudarzi I, Masoudi R, Ahmadi A, Momeni A. Effect of progressive muscle relaxation and aerobic exercise on anxiety, sleep quality, and fatigue in patients with chronic renal failure undergoing hemodialysis. *Int J Pharm Clin Res*. 2016; 8(12): 1634–9.
- [43] Paluchamy T, Vaidyanathan R. Effectiveness of intradialytic exercise on dialysis adequacy, physiological parameters, biochemical markers and quality of life - A pilot study. *Saudi J Kidney Dis Transpl*. 2018; 29(4): 902–10.
- [44] Jolly SE, Navaneethan SD, Schold JD, Arrigain S, Konig V, Burrucker YK, et al. Development of a chronic kidney disease patient navigator program. *BMC Nephrol [Internet]*. 2015; 16(1): 1–8.
- [45] D'souza B, Prabhu R, Unnikrishnan B, Kamath R. Effect of Multidimensional Educational Interventions Among Dialysis Patients. *Open Urol Nephrol J*. 2018;11(1):72–8.
- [46] Wakeel J Al, Harbi A Al, Bayoumi M, Al-Suwaida K, Ghonaim M Al, Mishkiry A. Quality of life in hemodialysis and peritoneal dialysis patients in Saudi Arabia. *Ann Saudi Med*. 2012; 32(6): 570–4.